TO ALL TO WHOM THESE: PRESENTS SHAME COME:

# The J. C. Robinson Seed Co.

Colhereus, there has been presented to the

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COLOR OF WHICH IS THE EURITON OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COLOR OF WHICH IS THE EURITON AND EXHIBITS, A COLOR OF WHICH IS THE EURITON AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE IN THE TIPLICANT (S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) FOR THE TERM OF EXPLICENT.

TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IN PRODUCING A HYBRID OR DIFFERENT ETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN

'CRIHT'

In Lestimony Wathereof, I have hereunto set my hand and caused the seal of the Elaut Tariety Protection Office to be affixed at the City of Washington day of March the year of our Lord one thousand nine hundred and eighty-five.

#### FURM APPROVED UNITED STATES DEPARTMENT OF AGRICULTURE QMB NO. 40-R3822 AGRICULTURAL MARKETING SERVICE LIVESTOCK, POULTRY, GRAIN & SEED DIVISION No certificate for plant variety protection may APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE pe is and unless a completed application form has been received (5 U.S.C. 553). INSTRUCTIONS: See Reverse, FOR OFFICIAL USE ONLY 1b. VARIETY NAME TEMPORARY DESIGNATION OF PV NUMBER VARIETY 8400042 CRLHt 1-2-2-1 XXX FILING DATE TIME 3. GENUS AND SPECIES NAME KIND NAME 2:30 1/17/84 P,M. Yellow Dent Corn Zea mays L. FEE RECEIVED DATE 1,000 1/17/84 B. DATE OF DETERMINATION FAMILY NAME (BOTANICAL) 500,00 January 1981 Gramineae TELEPHONE AREA 7. ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP NAME OF APPLICANT(S) CODE AND NUMBE 402-779-2531 68069 Waterloo, NE THE J. C. ROBINSON SEED CO. 11. DATE OF INCOR-10. IF INCORPORATED, GIVE STATE AND IF THE NAMED APPLICANT IS NOT A PERSON, FORM OF PORATION DATE OF INCORPORATION ORGANIZATION: (Corporation, partnership, association, etc.) 1904 Nebraska Corporation NAME AND MAILING ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERVE IN THIS APPLICATION AND RECEIVE ALL PAPERS: Dr. Frank D. Cloninger THE J.C. ROBINSON SEED CO. CHECK BOX BELOW FOR EACH ATTACHMENT SUBMITTED: 13A. Exhibit A, Origin and Breeding History of the Variety (See Section 52 of the Plant Variety Protection Act.) X 13B. Exhibit B, Novelty Statement. 13C. Exhibit C, Objective Description of the Variety (Request form from Plant Variety Protection Office.) X 13D. Exhibit D, Additional Description of the Variety. 144. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFI X NO SEED? (See Section 83(a). (If "Yes," answer 14B and 14C below.) YES DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE 14c. IF "YES," TO 14B, HOW MANY GENERATIONS OF PRODUC-TION BEYOND BREEDER SEED? LIMITED AS TO NUMBER OF GENERATIONS? REGISTERED CERTIFIED **FOUNDATION** YES X NO X NO (If "Yes," giv DID THE APPLICANT(S) FILE FOR PROTECTION OF THIS VARIETY IN OTHER COUNTRIES? name of countries and dates.) X NO (If "Yes," give name of countries HAVE RIGHTS BEEN GRANTED THIS VARIETY IN OTHER COUNTRIES? and dates.) DOES THE APPLICANT(8) AGREE TO THE PUBLICATION OF HIS/HER (THEIR) NAME(S) AND ADDRESS IN THE OFFICIAL 16. YES INO The applicant(s) declare(s) that a viable sample of basic seed of this variety will be furnished with the application and will be 17. replenished upon request in accordance with such regulations as may be applicable. The undersigned applicant(s) is (are) the owner(s) of this sexually reproduced novel plant variety, and believe(s) that the variety is distinct, uniform, and stable as required in Section 41, and is entitled to protection under the provisions of Section 42 of the Plant Variety Act. Applicant(s) is (are) informed that false representation herein can jeopardize protection and result in penalties. J. C. ROBINSON SEED CO DEC. 29, 1983 (SIGNATURE OF APPLICANT) (DATE) (SIGNATURE OF APPLICANT) (DATE)

FORM GR-470 (1-78)

FINANCE

VICE PRES -

CRIHt was selected from (Wll7Ht x Mol7Ht) a cross of an early Wisconsin inbred and a full-season Missouri released inbred line. This line was developed by self-pollination and selection for early silking and agronomic type for six generations. At Waterloo, NE, CRIHt flowers four days earlier than A632. This inbred typically produces a moderate height plant with ear attachment in the lower half of the plant. Plants have moderate length, moderate width, upright leaves and moderate length ears.

#### Corn Application No. 8400042

13A. Four generations of uniformity and stability have been observed. Slight plant height variability was observed. (186-206 cm. plant height with 55-75 cm. ear height in one sampling of 100 plants.) Phenotype was very uniform from plant to plant. CR1Ht ears generally have straight rows for about two-thirds of the ear length and the tip in many cases will have irregular rows. A small number of ears with straight rows are found.

13B. CR1Ht characteristics are more like Mo17Ht than any other common variety. However, flowering is earlier in CR1Ht.

### DAYS FROM PLANTING

	50% POLLEN			50% SILK		
	1981	1982	1983	1981	1982	1983
CR1Ht	69	76	79	70	77	79
Mo17	75	85	87	77	87	88

CR1Ht color is approximately one value darker on the Munsell color chart. The field sample we used of CR1Ht rated G4/4 and Mo17 rated G5/4. The leaf angle of CR1Ht is, in general, approximately  $10^{\circ}$  more upright than is Mo17Ht.

FDC/mrf 7/13/84

EXHIBIT (Corn)

#### UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION

HYATTSVILLE, MARYLAND 20782

# **OBJECTIVE DESCRIPTION OF VARIETY**

CORN (ZEA MAYS)			
NAME OF APPLICANT(S)	FOR OFFICIAL USE ONLY		
THE J. C. ROBINSON SEED CO.  ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	8400042		
ADDITION TO THE SECOND ONLY ONLY ONLY ONLY	VARIETY NAME OR TEMPORARY		
Waterloo, NE 68069	CR1HT RfS		
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (e.s. 0 8 9 or 0 9 ) when number is either 99 or less or			
1. TYPE:			
2 1 = SWEET 2 = DENT 3 = FLINT 4 = FLOUR 5 = PC	OP 6 = ORNA. ENTAL		
2. REGION WHERE BEST ADAPTED IN THE U.S.A.:			
2 1 = NORTHWEST 2 = NORTHCENTRAL 3 = NORTHEAST 5 = SOUTHCENTRAL 6 = SOUTHWEST 7 = MOST REGIONS	4 = SOUTHEAST		
	comments" (pg. 3) state how		
4 8 DAYS FROM EMERGENCE TO 50% OF PLANTS IN SILK	s were calculated) 7 8 HEAT UNITS		
DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY	HEAT UNITS		
7 2 DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 1 5	7 6 HEAT UNITS		
4. PLANT:			
1 9 2 CM. HEIGHT (To tessel tip)	6 0 CM. EAR HEIGHT (To base of		
THE CONTENTED CM. LENGTH OF TOP EAR INTERNODE	in the second of		
Number of Tillers: Number of Ears Per Stalk:			
and the second s	SLIGHT TWO-EAR TENDENCY		
Cytoplasm Type:	EAR TENDENCY 4 = THREE-EAR TEN		
Cytopiasiii 1 ypo.			
1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHER	(Specify)		
5. LEAF (Field Corn Inbred Examples Given):			
Color: 17. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1.			
2 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GRE	EN (B14) 4 = VERY DARK GREE		
Angle from Stalk (Upper half): Sheath Pubscence:			
3 $1 = < 30^{\circ}$ $2 = 30-60^{\circ}$ $3 = > 60^{\circ}$ 1 = LIGHT (			
Marginal Waves: Longitudinal Creases:			
3 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSENT			
Width: 3 = MANY (	PA11)		
0 9 CM, WIDEST POINT OF EAR NODE LEAF 0 7 0 CM. EA	AR NODE LEAF		

8. KERNEL (Dried):						
5 = BROW	RLESS 2 = RED-WH N 6 = LIGHT R EGATED (Describe)	ED 7.9	FAN 4 = BRONZ CHERRY RED			
RJS 12/14/84		EGATING (Describe)	5 = BRONZE	6 = RED		
3 1 WHITE 2 PIN 7 = PURPLE 8 = PAI	LE PURPLE 9 = VAR	-		· · · · · · · · · · · · · · · · · · ·		
2 Endosperm Color: 1 = WH	HITE 2 = PALE YELLOW	3 = YELLOW 4 =	PINK-ORANGE 5=	WHITE CAP,		
Endosperm Type:			• • •			
1 = SWEET (su1)	2 = EXTRA SWEET (sh2) 6 = HIGH PROTEIN	3 = NORMAL STARCH 7 = HIGH LYSINE	CH 4 = HIGH AMYLOSE STARCH 8 = OTHER (Specify)			
2 4 GM. WEIGHT /100 SEEDS (U	GM, WEIGHT /100 SEEDS (Unsized Sample)					
9. COB:  2 1 MM, DIAMETER AT MID-POI	NT					
Strength:  1 = WEAK 2 = STRO	· .	lor:  1 = WHITE 2 = PIN 5 = VARIEGATED	K 3 = RED 4 = BR 6 OTHER (Specify)			
10. DISEASE RESISTANCE (O = Not Tester	d, 1 = Susceptible, 2 = Resistant):					
O STALK ROT (Diplodia)  2 NORTHERN LEAF BLIGHT  O SOUTHERN RUST  O BACTERIAL LEAF BLIGHT	O STALK ROT (F O SOUTHERN LE O CORN SMUT	AF BLIGHT	O STALK ROT (GIR			
OTHER (Specify)	<b>L</b>		<b></b>	• ,		
11. INSECT RESISTANCT (O = Not Tested,	1 = Susceptible, 2 = Resistant):					
CORNBORER  O ROOTWORM (Northern)	EARWORM  ROOTWORM (Western)	О SAPBEE	rLE 0	APHID		
O ROOTWORM (Southern)	OTHER (Specify)					
12. VARIETIES MOST CLOSELY RESEMB	LINO THAT CHOMITTED FOR	THE CHARACTERS GIVEN	1:			
	VARIETY	CHARACTER	VARIE	TY		
CHARACTER	H99	Kernel Type	W117			
Maturity Plant Type	Mol7	Quality (Edible)				
Ear Type	Mol7	Usage	W117			
REFERENCES: U.S. Department Agriculture. Corn: Culture, Processing, Pro Emerson, R.A., G.W. Beadle, a	Yearbook 1937. ducts. 1970 Avi Publishing Com nd A.C. Fraser. A Summary of Li Crop Science Society of America	pany, Westport, Connecticut nkage Studies in Maize.Corne	(Numerous (Authors) H A.E.S., Mem. 180. 1935.			
Stringfield, G.H. Maize Inbred	Lines of Ohio, Ohio A.E.S. Bul.  of for the Classification of Corn In	831. 1959.	nio State University.	<b>)</b>		

## 13 D.

CRIHt ear type has the cob characteristics of Mol7. For example, when broken there are silks running through the interior tubes of the cob. CRIHt ears generally have straight rows for about two-thirds of the ear length and the tip in many cases will have irregular rows.